

10/570056

1AP20 Rec'd PCT/PTO 23 FEB 2006

SOLID STICK INSECTICIDAL COMPOSITION

Technical Field

The present invention relates to insecticidal compositions, and more specifically to both novel forms of insecticidal compositions and methods of controlling crawling pests such as cockroaches.

Background Discussion and Prior Art

10 Insects such as cockroaches and ants are a continual problem in both domestic, commercial and industrial establishments. They are difficult to control, finding harbourage in a multitude of locations including behind and under cabinets, appliances, sinks and worktops and in
15 cupboards. Furthermore, they can readily move to a new home when a harbourage becomes unacceptable.

Conventional methods for controlling such pests include aerosol sprays and baits. However, each of these methods has disadvantages. Aerosol products do not
20 generally have a long residual activity and the spray tends to radiate widely, going onto surfaces where protection is not sought or where the active ingredient is not wanted. In addition, aerosol propellants tend to be flammable.

25 Bait compositions come in a variety of forms including tablets, pastes, gels and other semi-solid preparations. Bait compositions are frequently held in a container or housing that only allows insect access. This means that baits cannot be placed in many areas where the
30 insects might walk or find harbourage, such as around the periphery of cabinet doors, on the undersides of worktops, or along junctions.

U.S. Patent No. 3,162,575 (Lang) discloses an insecticide stick for flies having 40-80%
35 microcrystalline wax base mixed with 10-60% petroleum oil, up to 50% particulate sugar and 0.25-1.5 dimethyl-

dichloro-vinyl-phosphate (DDVP) insecticide. The insecticide is applied to the faces of domestic livestock by rubbing, preferably between the eyes. The sugar reportedly causes the flies to be drawn toward the area of the face which is coated with the active ingredient.

U.S. Patent No. 3,826,232 (Duffey et al.) discloses a pest control stick for application to the neck or face of domestic animals to control fleas, lice, ticks, flies and the like. The stick comprises 0-(2-isopropoxyphenyl)-N-methyl carbamate as the active insecticide, fatty acid, fatty alcohol, polyethylene glycol and, preferably, a bitter tasting compound which serves to discourage licking by other animals and tasting by children.

U.S. Patent No. 4,473,582 (Greene) discloses an insecticidal stick for application to a household surface or domestic animals. The sticks comprise 0.25-5 wt% water-insoluble insecticide, 10-65 wt% fatty hydrocarbon monoether or propylene glycol and 30-70 wt% monoethanolamide of a fatty acid. When applied to a household surface, the stick reportedly deposits a thin film of the stick formulation on the surface and insects, such as cockroaches, walking or crawling over the film, it is believed, ingest or absorb the insecticide through their cuticles.

GB 608,715 (Ash Laboratories Limited) discloses solid insect control materials formed by compacting and subsequently drying a mixture of active insecticide, a liquid and a solid diluent or filler. The mixture may also contain a waxy or fatty substance so as to form a waxy crayon-like solid.

The solid insecticidal sticks described in the prior art address only some of the problems associated with the use of conventional pest control methods. Hence, there is still a clear need for improved compositions and methods for controlling pests, especially crawling insects such as cockroaches.

Summary of the Invention

Accordingly, this invention provides a solid stick insecticidal composition comprising a base, an insecticide
5 and at least 60 wt% of a food material.

This invention also provides a method for controlling insects in which a solid stick composition comprising an insecticide and at least 60 wt% of food material is applied to a hard surface to be treated.

10 Furthermore, this invention provides an insect control product comprising a solid stick insecticidal composition according to the invention which is packaged.

In addition, this invention provides for use of an insecticide and a food material in the manufacture a solid
15 stick insecticidal composition for application to a hard surface to be treated.

The compositions and methods of pest control of the present invention are highly effective against crawling insects and provide significant improvements over the
20 insecticidal sticks of the prior art. In particular, the compositions of the present invention provide significant advantages over known stick compositions when applied to hard surfaces. The high percentage of food material in the stick compositions results in the majority of the
25 deposited material being ingested by insects, such as cockroaches, thus leaving little or no material to be cleared away. In contrast, the prior art sticks have a tendency to remain in situ once deposited and have to be cleaned up. The insecticidal sticks of the present
30 invention are therefore considerably easier and more convenient to use than those described previously. Furthermore, less material is required to be deposited to obtain equivalent levels of pest control, e.g. kill rates. The insecticidal sticks of the present invention are also
35 stable and possess a long residual activity.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in this specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

Detailed Description of the Invention

As used herein, all figures given as %wt are, unless specified otherwise, the percentage by weight of that component relative to the total weight of the solid stick insecticidal composition. Where a set of parameter ranges are given for a particular component or composition, it will be understood that any lower limit in that set of ranges may be combined with any upper limit in the same set of ranges to provide a suitable range limit for that component or composition.

Where an ingredient can perform more than one function, for example act as both a food material component and as part of the base composition, it will be understood that, unless specifically indicated otherwise, it can contribute to the stated weight ranges relating to both functions equally.

The compositions of the present invention comprise at least one insecticide component. Suitable insecticides may be chosen from a wide range of active ingredients, both natural and synthetic.

Examples of suitable insecticide ingredients include pyrethroids, neonicotinoids (e.g. imidacloprid, thiamethoxam), avermectins, spinosyns (e.g. spinosad), hydramethylnon, fluorinated sulfluoramides, organophosphates including diazinon and chlorpyrifos, pyrazoles such as fipronil, chlorfenapyr, indoxacarb, borates, benzoylphenyl ureas, carbamates and hydrazones. Preferred insecticides include organophosphates, such as chlorpyrifos, pyrazoles, such as fipronil, and indoxacarb.

One or more insecticides may be employed. In addition, insecticide(s) may be micro-encapsulated, non-encapsulated or a combination of the two. For example, a combination of non-encapsulated and microencapsulated chlorpyrifos may be used. The particular choice of insecticide(s) depends on several factors including the target insect(s), the strategy for killing the insects, the regulatory approval status in a particular country, cost, etc. As regards the strategy for killing insects, insecticides can act in a variety of ways. For example, some insecticides directly kill the insects, some affect the fecundity of the insect such that the insect population is reduced in subsequent generations, and some alter the behaviour of the insect in a manner that will bring about its destruction. In addition, insect populations can also be reduced as a result of transmission of insecticidal composition between insects - this is generally referred to as "secondary kill".

The total amount of insecticide in the solid stick insecticidal composition will suitably range from 0.01 to 40 wt%, preferably from 0.01 to 20 wt%, more preferably from 0.01 to 10 wt%, yet more preferably from 0.01 to 5 wt%, even more preferably from 0.02 to 5 wt%, yet more preferably from 0.03 to 3 wt% and most preferably from 0.03 to 2 wt%.

Biological control agents, such as the bacterium *Bacillus thuringiensis* or entomophagous fungi (e.g.

Beauveria bassiana) could also potentially be used as 'active ingredients' in the present invention. For the purposes of the present invention, any amount of biological control agent present is considered to form
5 part of the insecticide component as regards weight ranges.

The compositions of the present invention comprise at least one food material component.

Examples of suitable food materials include
10 carbohydrates, proteins, lipids, essential oils, water, spices and mixtures thereof. Preferred carbohydrates include honey, sugars (including monosaccharides and disaccharides), oligosaccharides, polysaccharides and other complex carbohydrates such as starches, pectins and
15 cellulosic materials. Examples of suitable starches include those found in flours such as, for example, corn flour and potato flour. Other preferred carbohydrate and/or protein food materials include soya flour, cereal products such as bran and wheat germ, vegetable products,
20 dehydrated vegetables, dried yeast, egg powder and insect derived materials such as silkworm pupa, eggs and body parts.

Other preferred food materials include fats and oils such as fully hydrogenated fatty acids. A preferred oil
25 is soya bean oil. Other fatty materials that may be used include esters, waxes, soaps, phospholipids such as lecithin, glycolipids, terpenes and steroids.

The total amount of food material in the solid stick insecticidal compositions is at least 60 wt%. Suitably,
30 the food material is present in the insecticidal composition in an amount of at least, in the order of preference, 65, 70, 75, 80, 85 and 90 wt% to 99.99 wt%. Particularly preferred ranges are from 70 to 99.99 wt%, more preferably from 90 to 99.99 wt%.

35 The insecticide is dispersed within, and the food material either forms part of and/or is dispersed within,

a fatty base. The base composition may also have dispersed therein one or more optional components. Preferably, all the components of the stick are uniformly dispersed within the base composition.

- 5 The base ingredients contribute to the physical and rheological properties of the stick. Some ingredients may contribute primarily to the physical properties of the stick, i.e. the hardness or rigidity of the stick, some may act primarily as lubricants, some may act primarily as
10 emollients or emulsifiers and some may act primarily as adhesion promoters or binding agents. It may also be beneficial to employ ingredients in the base which act primarily as solvents for certain components, for example certain insecticide components, of the insecticidal stick.
- 15 A preferred insecticide of the present invention is chlorpyrifos and this is preferably dissolved in a suitable solvent component of the base composition such as, for example, hydrogenated palm stearine and soya bean oil.
- 20 Suitable base ingredients include fatty alcohols, fatty acids, various esters including lower alkyl esters of fatty acids, polyethylene glycols (carbowaxes), monoalkanolamides, castor oil, mineral oils and petroleum jellies. Suitable base materials which contribute to the
25 hardness and rigidity of the sticks are typically materials having higher melting points. Such materials are also important in the moulding process. These materials are typically waxes or wax-like materials and include carnauba wax, candelilla wax, amorphous
30 hydrocarbon waxes, petroleum-based waxes such as microcrystalline wax, beeswax, paraffin waxes, cocoa butter, hydrogenated vegetable oils such as hydrogenated palm stearine and hydrogenated castor oil, lanolin and lanolin absorption bases, some paraffin oils, lecithin and
35 silicone waxes.

Some ingredients can have a dual function as a food material and as a component of the base. Examples of such dual functionality ingredients include hydrogenated palm stearine, tallow, lard, suet and duck fat. In addition, 5 some food materials that are not at all waxy or fatty in nature also act as base ingredients. For example, flours, e.g. soya flour, can act as binding agents and thus effect the rheology of the base composition. In addition, ingredients such as lecithin and monoglycerides can act as 10 emulsifiers, helping to suspend and disperse hydrophilic materials in hydrophobic materials and vice-versa. Preferably, the base composition comprises one or more materials which act as emulsifying agents. Such emulsifying materials are suitably present in amount of 15 from 1 to 10 wt%, preferably from 2 to 8 wt% of the solid stick insecticidal composition.

Preferably, the base comprises at least one of hydrogenated palm stearine, soya flour, lecithin and monoglyceride.

20 Suitably, the base composition has a melting point in the range from 20 to 200 °C, preferably from 35 to 100 °C and more preferably from 40 to 80 °C.

The total amount of base ingredients in the solid stick insecticidal composition will suitably be, in order 25 of preference, at least 10, 20, 30 and 40 wt%. Suitably, the total amount of base ingredients in the solid stick insecticidal compositions will be, in order of preference, no more than 99.99, 99, 95, 90, 85, 80, 75 and 70 wt%.

The insecticidal compositions may also comprise one 30 or more optional components such as attractants, preservatives, anti-oxidants, feeding stimulants, fillers, animal (including human) taste deterrents and colorants.

In a preferred embodiment, the insecticidal compositions comprise at least one attractant. Where 35 compositions comprise a food material and an attractant, this may be achieved by using separate food material and

attractant components or alternatively by using an ingredient that acts both a food material and an attractant.

Examples of suitable attractants, which attract
5 through smell and/or other neurosensory pathways, include pheromones, yeast extracts, complex foods and volatile materials such as flavour essences and food derivatives including fenugreek, maple lactone, herbs and spices. When insecticides are themselves repellent in nature to
10 insects, an attractant as taught and disclosed in WO 97/00610 (R & C Products Pty Ltd) may suitably be used. Preferred attractants include maple lactone, soya bean oil, malt extract and honey.

Examples of dual functionality food materials, which
15 also act as attractants, include soya bean oil, spices, yeasts, yeast extracts, malt extracts and maple lactone. The function of each material will of course depend to some extent on the particular species of insects you are looking to treat. For example, soya bean oil, honey, malt
20 extract and maple lactone all act both as food materials and attractants for cockroaches. Preferably, the insecticidal compositions of the present invention comprise maple lactone, which is suitably present in an amount of from 0.001 to 1 wt%, preferably from 0.005 to
25 0.5 wt % and more preferably from 0.005 to 0.05 wt%.

Suitably, the total amount of attractant in the insecticidal compositions (including any dual
functionality components) is at least 0.1, preferably at least 0.5, more preferably at least 1.0 and yet more
30 preferably at least 2 wt%.

In contrast to the prior art stick compositions which are only effective when carefully applied as an unbroken line such that the insects invariably cross the line at some point in their travels, the insecticidal compositions
35 of the present invention can be applied to hard surfaces as "spots" or short lines of material whilst still

providing a level of pest control that is at least as equivalent and typically more effective than that described in the prior art. This is particularly the case when the compositions contain an attractant.

5 A preservative may be present to inhibit the growth of microorganisms on the composition, as such growth may potentially repel insects. Amongst the preservatives that may be used are sorbic acid and salts thereof (e.g. potassium sorbate), DowicilTM (Dow-Elanco) and methyl- and
10 propylparabens. Suitably, preservatives may be present in an amount from 0.0001 to 10 wt%, preferably from 0.001 to 5 wt%.

Anti-oxidants may be present to maintain food material palatability in an amount of from 0.0001 to 10
15 wt%, preferably from 0.001 to 5 wt%. Suitably, in compositions that include fats and/or oils, an antioxidant such as tert-butyl hydro-quinone (TBHQ), butylated hydroxytoluene or butylated hydroxyanisole is used. TBHQ is a preferred anti-oxidant in insecticidal sticks of the
20 present invention.

Animal taste deterrents may be included to minimise the risk of accidental consumption. Suitable human deterrents include denatonium benzoate (BitrexTM MSL). When present, a human taste deterrent is suitably used in
25 an amount of from 1 to 200 ppm, preferably from 1 to 100 ppm, more preferably from 10 to 50 ppm and most preferably from 5 to 20 ppm.

The sticks may be coloured by including one or more colorants in the formulation. Suitable colorants include,
30 for example, many of the conventional food dyes, e.g. carmoisine. Suitably, a colorant may be present in an amount of up to 1 wt%, preferably from 0.0005 to 0.5 wt%, more preferably from 0.0005 to 0.2 wt% and yet more preferably from 0.0005 to 0.05 wt%.

35 The sticks of the present invention leave a deposit of composition on a hard surface when applied to the hard

surface. The sticks should not bend, crumble, crack or break during application. The physical properties and quality of the stick during manufacture, storage and use are determined for the most part by the composition of the fatty base composition. These properties and qualities are largely related to the rheology of the mixture at various temperatures. For example, during manufacture (usually while warm), it is preferably possible to mix or mill the formulation and to pour and mould it whilst holding any insoluble components evenly dispersed without settling. During shelf life and the life of usage, the stick should remain rigid and stable. In use, the stick must be capable of depositing composition onto hard surfaces on application of a downward force. The hard surfaces may, for example, be tiles, concrete, wood, stone, metal, glass or plastics such as melamine. It will be understood that hardness and deposition properties of the solid insecticidal sticks will vary somewhat depending on the composition, and in particular the base composition. The stick may be relatively hard and akin to a conventional drawing crayon. In such cases, the stick may have to be drawn across, rubbed onto or, where spot deposition is required, twisted onto the hard surface in order to deposit insecticidal composition. Alternatively, the stick may be "softer" and more akin to a lipstick. Clearly more force is ordinarily required to deposit a drawing crayon than a lipstick onto a hard surface. Softer compositions will also tend to deposit smears of material. It will also be evident that depending on the nature of the surface to be treated, different formulations may be preferred in order to achieve satisfactory deposition. A crayon-like stick will readily deposit on rough surfaces such as concrete whereas it may have difficulties depositing on shiny surfaces such as stainless steel or polished stone, where a softer more adherent formulation may be more appropriate.

The deposited composition suitably adheres to the hard surface, i.e. once applied to the surface, and remains more or less in place until ingested by insects or physically wiped or scraped off. Suitably the
5 compositions do not dry out to any significant degree after application (at least over a period of a day, and preferably for longer) and the deposited material remains in a form which allows it to be ingested by the target insects and therefore does not harden to a point where it
10 is unpalatable. In this way, insects are encouraged to approach the deposited composition and/or remain with the deposited composition for a period of time.

The solid sticks of the present invention suitably have the outward appearance of a cylinder, rod or crayon.
15 The sticks may be any suitable shape in cross section including circular or a polygon of three or more sides. A preferred shape is circular. The sticks may be tapered at one or both ends, preferably one end.

The sticks are typically of a size suitable for
20 holding in the hand so as to allow the easy drawing of lines or spots on hard surfaces. Suitably, the sticks are from 10 to 300 mm in length, preferably from 30 to 200 mm, and more preferably from 50 to 120 mm. Suitably, the average cross-sectional area of the stick is from 20 to
25 2,000 mm², preferably from 40 to 500 mm² and more preferably from 60 to 200 mm².

The hardness of the stick should be such that the crayon can be held near the surface to be treated and pressed onto and dragged across that surface with a
30 moderate amount of force without breaking. At the same time it should be capable of depositing a sufficient quantity of material for insects such as cockroaches to ingest a lethal dose.

The hardness and rigidity of the solid stick
35 compositions may be determined by penetrometry. Suitable procedures are described in ASTM D1321; IP376; DIN 51579.

For example, hardness may be measured using a lab plant PNT penetrometer equipped with a Seta wax needle (weight 2.5 grams) which has a cone angle at the point of the needle specified to be $9^{\circ}10'$ \pm $15'$. The barrel of the stick is cut to leave a flat uniform surface. The needle is lowered onto the surface of the composition and then a penetration hardness measurement is conducted by allowing the needle with its holder to drop under a total weight, (ie. the combined weight of needle and holder) of 50 grams for a period of five seconds after which the depth of penetration is noted. Desirably the test is carried out at a number of points, e.g. six points, on each sample and the results are averaged.

Utilising a test of this nature, the solid insecticidal compositions of the present invention suitably have a penetration hardness of from 0.1 to 10 mm, preferably from 0.5 to 8 mm, more preferably from 1 to 5 mm, yet more preferably from 1 to 4 mm and even more preferably from 1 to 3 mm.

The solid stick compositions may also be tested for their ability to deposit onto hard surfaces. For example, a suitable deposition test is to fit a sample of the composition with standardised shape and size to an apparatus which draws the sample across a test surface under standardised conditions. The amount transferred to the surface is then determined as an increase in the weight of the substrate to which it is applied. Suitable test substrates include ceramic tiles, concrete, wood, melamine (laminated), metal and glass.

The substrates are weighed before use. The sticks are previously unused. The apparatus comprises a flat base to which a flat substrate is attached by a clip at each end. A pillar having a mounting to receive a standard size stick is mounted on an arm that is moveable horizontally across the substrate by means of a pneumatic piston. Each stick is kept at ambient laboratory

temperature overnight before the measurement is made. The stick is placed in the apparatus and a spring positioned to bias the stick against the substrate with a standardised force. The apparatus is operated to pass the
5 stick laterally across the substrate a set number of times. The substrate is carefully removed from the rig and reweighed.

Such a deposition test may also be used to help assess the quality and nature of the deposition, e.g. the
10 adhesion properties, whether smearing is occurring and stability properties (e.g. drying-out properties).

Insecticidal compositions are often widely used in warm countries. For this reason, it is preferable that the stick compositions of the present invention have a
15 reasonably high melting point. Suitably, the solid stick insecticidal compositions have a melting point of at least 20 °C, preferably at least 35 °C, more preferably at least 40 °C and even more preferably at least 45 °C.

The solid stick insecticidal compositions preferably
20 have a setting point in the range from 20 to 200°C, more preferably from 30 to 70°C and yet more preferably from 45-55 °C.

The stick may be used in an uncovered form. Alternatively, and preferably, the stick may be provided
25 with a packaging. The packaging may cover the entire length of the stick or only part thereof. It may comprise of a protective wrapping (paper, plastic, fabric, metal foil or other suitable material including laminated materials) axially orientated around the stick, which is
30 suitably affixed, e.g. by gluing, along a line of overlap or as a self-adhesive sticker wrapping. The wrapping may be removed entirely before use of the stick. However, a preferred embodiment of this invention employs wrappings which can be readily torn to reveal "fresh" stick.
35 Preferably such wrappings are designed to allow progressive tearing at predetermined points. This may be

achieved, for example, by using an excess of wrapping material to generate a flap of wrapping material beyond the line of overlap and fixing. Tear lines or points of weakness are then generated at regular intervals along the flap as starting points for tearing off strips of wrapping. For example, serrations can provide points of weakness, especially in plastic wrappings. Other suitable tear starting points include short cuts and perforated lines in the flap of wrapping material.

10 Alternatively, the packaging may comprise an applicator made from, for example, plastic or metal. A simple applicator construct may comprise a bottom, in which is held one end of the stick, and a tubular lid, which may partially or fully enclose the insecticidal stick. In order to apply the insecticidal composition, the lid is removed and then replaced after use. More complex applicators enable the insecticidal stick to be extended from a casing. This may be achieved by having the insecticidal stick on a threaded screw and rotating the screw by turning the bottom of the applicator. Alternatively, the stick may be extended from and retracted into a casing by sliding it within the casing. Typically this may be achieved by actuating an arm within a slot in and extending from the casing, the arm being connected to a housing holding the stick within the casing. The art is replete with examples of suitable applicators which may be employed with the insecticidal stick of the present invention. For example, applicators used in personal care for delivery of lipsticks, lipsalves and deodorants, are particularly useful.

The packaging itself can comprise a label or alternatively a label can be applied to the packaging. Suitably, the label includes information on the product, such as the brand name etc., and instructions for use. A major advantage of using a packaging is that it reduces the level of direct human contact with the insecticidal

composition. In addition, wrappings tend to increase the strength of the sticks. Packagings in which the sticks are enclosed within rigid casings allow for stick compositions to be used that may not be entirely suitably for use as "crayon-like" products because, for example, they are softer in nature. Preferably, the packaging material is such that it does not allow the insecticidal composition, or components thereof, to leach into or out through it. Suitable wrapping materials in this respect include paper, plastic films (e.g. polypropylene), fabric, metal foil and laminates of these. When using paper or fabric as wrapping materials, it is desirable to coat the side of the wrapping material coming into contact with the insecticidal composition with a suitable protective layer, such as an oil resistant layer. Examples of suitable paper wrappings include wax-coated paper, polymer film coated paper (e.g. milk carton materials) and paper which has been solvent or chemically treated, such as siliconised paper or greaseproof paper.

The sticks can be manufactured using techniques already known in the art. For example, the methods used to manufacture personal care products such as lipsticks and lipsalves can be readily imported and adapted for the manufacture of the insecticidal sticks of the present invention. Such methods are well-known and described, for example, in Harry's Cosmeticology, (1982), 7th Edition, Longman Scientific & Technical, U.K. In addition, the techniques employed in the manufacture of waxy crayons can also be imported and adapted for the manufacture of the present stick compositions. All of the aforementioned manufacturing processes typically employ melt cast processes in which the formulation is prepared as a molten mixture, poured into moulds, allowed to set and finally removed. Accordingly, the insecticidal sticks of the present invention may be prepared by :

- (i) forming a molten, pourable mixture of the insecticidal composition;
- (ii) pouring the molten mixture into moulds;
- (iii) allowing the mixture to set in the moulds; and
- 5 (iv) releasing the solid stick insecticidal compositions from the moulds.

In order to facilitate setting and good surface properties, the moulds may be cooled either prior to and/or during step (iii). This may, for example, be
10 achieved through circulating water, which itself may have been chilled. The moulds are suitably made from metal, for example brass, and are commonly referred to as pipe moulds. The molten composition is poured onto a flatbed moulding table containing multiple holes (often
15 thousands), each hole being suitably shaped to form, for example, a stick with a circular cross section and having a tapered end. As the molten compositions settles into the moulds, it is optionally cooled by circulating water. Once the insecticidal sticks have cooled and solidified,
20 any excess material is scraped away and the sticks are removed from the moulds. Removal may be achieved by pushing rods, which sit at the base of the moulds during the casting process, into the moulds. The sticks may be further shaped once released. For example, they may be
25 shaved or cut into appropriate lengths. The sticks may then go on to be packaged as it appropriate.

It is also possible to produce the insecticidal compositions of the present invention by extrusion techniques.

30 The insecticidal sticks of the present invention may be used to control any insects which tend to land and/or crawl over hard surfaces such as ants, cockroaches, silverfish, terrestrial crustaceans such as woodlice (slaters), stored product pests, and flies. The
35 insecticidal stick compositions are particularly effective at controlling insects which crawl over hard surfaces and

have been found to be particularly effective at controlling the *Blattella germanica* (German cockroach) and *Periplaneta americana* (American cockroach).

The invention will now be further illustrated with
5 reference to the following non-limiting examples:

EXAMPLESExample 15 Solid stick insecticidal formulation

<u>Component</u>	<u>Wt%</u>	<u>Function</u>	
Palm Stearine Hydrogenated	32.865	Base	
10 component/food material			
Soya bean Oil	7.455	Food	
material/attractant			
Soya Lecithin	4.970	Base	component/food
material			
15 Distilled Monoglyceride	1.491	Base	component/food
material			
Malt Liquid Extract Ltnd	12.921	Food	
material/attractant			
Honey Manufacturing Grade	24.849	Food	
20 material/attractant			
Soya Flour Full Fat	13.945	Base	component/food
material			
Chlorpyrifos 99%	0.353	Insecticide	
Chlorpyrifos (MC) (#)	0.753	Insecticide	
25 Denatonium Benzoate	2.5 % 0.040	Human	taste
deterrent			
Maple Lactone	0.010	Food	
material/attractant			
Potassium Sorbate	0.298	Preservative	
30 TBHQ	0.050	Anti-oxidant	

Chlorpyrifos (MC) is a microencapsulated chlorpyrifos, chlorpyrifos being present at 20 wt% of the capsule weight.

35

Method of preparation

The formulation given above was prepared as follows:

- i) With the temperature set to 70 °C the soy bean oil and palm stearine were added to a mixer and stirred.
- 5 ii) Stirring was continued for 20 minutes, the temperature adjusted to 60°C and the potassium sorbate added.
- iii) Stirring was continued until the materials were well blended and temperature was stable at 60 °C.
- 10 iv) The soy lecithin was preheated to 30 °C and added to the mixture.
- v) Stirring was continued until the mixture was well blended and then the distilled monoglyceride was slowly added.
- 15 vi) The malt and honey were preheated 30 °C and then slowly added, with the stirrer speed reduced, along with the benatonium benzoate.
- vii) The mixture was stirred until well blended
- 20 viii) The Tenox TBHQ was added to the mixture and stirring continued until well blended - some needles may still be visible at this stage
- ix) The soy flour was slowly added through a sieve and the mixture stirred until no lumps were visible and the mixture was golden yellow and smooth.
- 25 x) Both the 99% chlorpyrifos and MC chlorpyrifos and maple lactone were added to the mixture and the mixture mixed until it was uniform.
- xi) Mixing was continued at temperature of 57 to 60 °C for 15 minutes
- 30 xii) A sample was taken for testing
- xiii) The tested mix was then poured into brass crayon moulds and cooled (in this case refrigerated below 0 °C.)
- xiv) Once hardened (after about 15 minutes) they were
35 pushed out of the brass moulds using a plastic plunger.

- xv) The crayons were wrapped in self-adhesive plastic wrappers with one end of wrapper flush with end of the stick, exposing approximately 10 mm of crayon at the other end.

5

The temperature of the mixture was never allowed to rise above 65 °C during either the mixing or filling stages.

Assay

10

The efficacy of the crayon composition was compared with that of a gel-type cockroach bait composition containing the same active ingredient in the same concentration. The gel-type bait was applied as "blobs" onto the plastic test
15 arenas. The crayon composition was drawn as smears onto the plastic test arenas.

- i) 8 mixed age and sex American cockroaches were placed in each of 6 plastic arenas (base area 1,785 cm² and
20 height 19 cm); 100 mixed age and sex German cockroaches were placed in each of 6 plastic arenas (base area 988 cm² and height 11cm). The arenas were covered with ventilated plastic lids to prevent escape. There were three replicates (arenas) for
25 each of the bait treatments.

- ii) The cockroaches were provided with a harbourage in the form of half an up-turned egg carton, a water source and a dog food pellet. The positioning of the harbourage, food and water was the same for each
30 arena.

- iii) The cockroaches were allowed to acclimatise overnight and any dead were replaced the next morning.

- iv) The compositions to be tested were applied as smears (crayon composition) or blobs (gel-type composition)
35 of approximately 0.5 g to the surface of small plastic trays and the plastic trays placed in the

same position in each of the treatment arenas. There were six arenas for each of the two treatments (3 American cockroaches, 3 German cockroaches). The compositions were left in each arena for a further three days.

- v) Cockroach mortality was assessed at the end of this period and the mean percentage mortality calculated.

Results

10

Species	% mortality after 3 days	
	Crayon	Gel comparison
American cockroaches	89	51
German cockroaches	67	62

It is evident from these results that the composition of the present invention demonstrated a superior efficacy against both species of cockroaches when compared to a conventional gel-type bait composition.

Example 2

The following solid stick insecticidal formulations were prepared:

5

Formulation 2a

	<u>Component</u>	<u>Wt%</u>	<u>Function</u>
	Palm Stearine Hydrogenated	32.965	Base component/food
10	material		
	Soya bean Oil	7.455	Food
	material/attractant		
	Soya Lecithin	4.970	Base component/food
	material		
15	Distilled Monoglyceride	1.491	Base component/food
	material		
	Malt Liquid Extract Ltnd	12.921	Food
	material/attractant		
	Honey Manufacturing Grade	24.849	Food
20	material/attractant		
	Soya Flour Full Fat	13.945	Base component/Food
	material		
	Chlorpyrifos 99%	0.506	Insecticide
	Denatonium Benzoate 2.5 %	0.040	Human taste deterrent
25	Maple Lactone	0.010	Food
	material/attractant		
	Potassium Sorbate	0.298	Preservative
	TBHQ	0.050	Anti-oxidant
	Carmoisine (1.5% w/w soln)	0.500	Colorant

30

Formulation 2b

	<u>Component</u>	<u>Wt%</u>	<u>Function</u>
	Palm Stearine Hydrogenated	33.465	Base component/food
35	material		

	Soya bean Oil	7.455	Food	
	material/attractant			
	Soya Lecithin	4.970	Base	component/food
	material			
5	Distilled Monoglyceride	1.491	Base	component/food
	material			
	Malt Liquid Extract Ltnd	12.921	Food	
	material/attractant			
	Honey Manufacturing Grade	24.849	Food	
10	material/attractant			
	Soya Flour Full Fat	13.945	Base	component/food
	material			
	Chlorpyrifos 99%	0.506	Insecticide	
	Denatonium Benzoate 2.5 %	0.040	Human taste deterrent	
15	Maple Lactone	0.010	Food	
	material/attractant			
	Potassium Sorbate	0.298	Preservative	
	TBHQ	0.050	Anti-oxidant	

Other suitable solid stick insecticidal formulations are:

Formulation 2c

5	<u>Component</u>	<u>Wt%</u>	<u>Function</u>
	Palm Stearine Hydrogenated	33.766-33.961	Base
	component/food material		
	Soya bean Oil	7.455	Food
	material/attractant		
10	Soya Lecithin	4.970	Base component/food
	Distilled Monoglyceride	1.491	Base component/food
	material		
	Malt Liquid Extract Ltnd	12.921	Food
	material/attractant		
15	Honey Manufacturing Grade	24.849	Food
	material/attractant		
	Soya Flour Full Fat	13.945	Base component/food
	material		
	Fipronil 97.5%	0.010-0.205	Insecticide
20	Denatonium Benzoate 2.5 %	0.040	Human taste deterrent
	Maple Lactone	0.010	Food
	material/attractant		
	Potassium Sorbate	0.298	Preservative
	TBHQ	0.050	Anti-oxidant

25

Formulation 2d

	<u>Component</u>	<u>Wt%</u>	<u>Function</u>
	Palm Stearine Hydrogenated	31.930-33.869	Base
30	component/food material		
	Soya bean Oil	7.455	Food
	material/attractant		
	Soya Lecithin	4.970	Base component/food
	material		
35	Distilled Monoglyceride	1.491	Base component/food
	material		

	Malt Liquid Extract Ltnd	12.921	Food
	material/attractant		
	Honey Manufacturing Grade	24.849	Food
	material/attractant		
5	Soya Flour Full Fat	13.945	Base component/food
	material		
	Indoxacarb 98%	0.102-2.041	Insecticide
	Denatonium Benzoate 2.5 %	0.040	Human taste deterrent
	Maple Lactone	0.010	Food
10	material/attractant		
	Potassium Sorbate	0.298	Preservative
	TBHQ	0.050	Anti-oxidant

Example 3

A solid stick of formulation 2b was prepared according to the method described in Example 1. The crayon had a diameter of 13 mm, a length of 78 mm and weighed 10 g.

For comparative purposes, a stick of the composition described in US 3,162,575 (Lang) was prepared. The Lang stick had the same dimensions as the stick of formula 2b and a similar weight. The stick comprised:

60 wt% microcrystalline wax - made up of a combination of white 1275/7 (ex Frank B Ross) and HiMic 1070 (ex BP Global Special Products) to give a melting point of 71 °C as required by Lang.

20 wt% medium viscosity oil.

20 wt% powdered sucrose.

The hardness of solid stick compositions were determined by penetrometry using a lab plant PNT penetrometer equipped with a Seta wax needle (weight 2.5 grams) and having a cone angle at the point of the needle specified to be 9°10' +/- 15'. The barrels of sticks were cut to leave a flat uniform surface. The needle was lowered onto the cut surface and a penetration hardness measurement conducted by allowing the needle with its holder to drop under a total weight, (ie. the combined weight of needle and holder) of 50 grams for a period of five seconds after which the depth of penetration was noted. The test was carried out at six points on the formula 2b stick sample and nine points on the Lang stick sample and the results averaged.

3 runs on the formula 2b stick gave average values of 2.52, 2.55 and 1.98 mm

1 run on the formula 2b stick gave an average value of 2.12 mm

1 run on the Lang stick gave an average value 5.59 mm.

5

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without
10 departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

CLAIMS

- 5 1. A solid stick insecticidal composition comprising a base, an insecticide and at least 60 wt% food material.
2. An insecticidal composition according to claim 1
10 comprising at least 65, preferably at least 70 and more preferably at least 75 wt% food material.
3. An insecticidal composition according to claim 1 or claim 2 comprising up to 99.99 wt% food material.
15
4. An insecticidal composition according to any preceding claim comprising food material selected from
20 (i) carbohydrates;
(ii) protein/carbohydrate containing materials;
(iii) fatty or oily materials; and mixtures thereof.
5. An insecticidal composition according to claim 4, in which the carbohydrates are selected from
25 (i) honey;
(ii) sugars including monosaccharides and dissaccharides;
(iii) polysaccharides and complex carbohydrates including cellulosic materials, pectin and starches; and mixtures thereof.

6. An insecticidal composition according to claim 4 or claim 5, in which the protein/carbohydrate containing materials are selected from:
- 5 (i) cereal products,
 - (ii) flour, including soya flour;
 - (iii) spices;
 - (iv) vegetable products including dehydrated vegetables;
 - (v) dried yeast;
 - 10 (vi) egg powder;
 - (vii) insect-derived materials including silkworm pupa, eggs and body parts; and mixtures thereof.
7. An insecticidal composition according to any one of
- 15 claims 4 to 6, in which the fatty or oily materials are selected from:
- (i) lipids;
 - (ii) essential oils;
 - (iii) fully hydrogenated fatty acids;
 - 20 (iv) vegetable oils;
 - (v) animal-derived fats;
 - (vi) esters;
 - (vii) waxes;
 - (viii) soaps;
 - 25 (ix) phospholipids;
 - (x) glycolipids;
 - (xi) terpenes;
 - (xii) steroids; and mixtures thereof.
- 30 8. An insecticidal composition according to any preceding claim comprising at least 10, preferably at least 20 and more preferably at least 30 wt% base ingredients.
- 35 9. An insecticidal composition according to any preceding claim comprising base ingredients selected

from fatty alcohols, fatty acids, various esters including lower alkyl esters of fatty acids, castor oil, mineral oils, petroleum jellies, polyethylene glycols, monoalkanolamides, carnauba wax, candelilla wax, amorphous hydrocarbon waxes, petroleum-based waxes including microcrystalline waxes, beeswax, paraffin waxes, cocoa butter, hydrogenated vegetable oils, lanolin, lanolin absorption bases, paraffin oils, lecithin, silicone waxes, flour, monoglycerides and animal derived-fats including tallow, lard, suet and duck fat.

10. An insecticidal composition according to any preceding claim comprising one or more dual functionality materials acting as both a food material and a base ingredient.
11. An insecticidal composition according to claim 10, in which the dual functionality material is selected from hydrogenated palm stearine, tallow, lard, suet, duck fat, monoglycerides, flours and lecithins.
12. An insecticidal composition according to any preceding claim in which the base comprises one or more emulsifying agents.
13. An insecticidal composition according to any preceding claim, in which the insecticide is selected from pyrethroids, neonicotinoids, avermectins, spinosyns, hydramethylnon, fluorinated sulfluoramides, organophosphates, pyrazoles, chlorfenapyr, indoxacarb, borates, benzoylphenyl ureas, carbamates, hydrazones and mixtures thereof.
14. An insecticidal composition according to claim 13, in which the insecticide comprises an organophosphate.

15. An insecticidal composition according to claim 14, in which the organophosphate is chlorpyrifos.
- 5 16. An insecticide according to claim 13, in which the insecticide comprises a pyrazole.
17. An insecticide according to claim 16, in which the pyrazole is fipronil.
- 10 18. An insecticide according to claim 13, in which the insecticide comprises an indoxacarb.
19. An insecticidal composition according to any preceding claim comprising insecticide in an amount of from 0.01 to 40 wt%, preferably from 0.01 to 20 wt% and more preferably from 0.01 to 10 wt%.
- 15 20. An insecticidal composition according to any preceding claim comprising an attractant.
- 20 21. An insecticidal composition according to claim 20, in which the food material also functions as an attractant.
- 25 22. An insecticidal composition according to claim 21, in which soya bean oil is a food material and an attractant.
- 30 23. An insecticidal composition according to claim 21 or claim 22, in which honey is a food material and an attractant.
- 35 24. An insecticidal composition according to any one of claims 21 to 23, in which malt extract is a food material and an attractant

25. An insecticidal composition according to any one of claims 21 to 24, in which maple lactone is a food material and an attractant.
- 5 26. An insecticidal composition according to claim 25, in which maple lactone is present in an amount of from 0.001 to 1 wt%, preferably from 0.005 to 0.5 wt %.
- 10 27. An insecticidal composition according to any one of claims 20 to 26 comprising an attractant which does not function as a food material.
- 15 28. An insecticidal composition according to any preceding claim further comprising a preservative.
- 20 29. An insecticidal composition according to claim 28, in which the preservative is present in an amount of from 0.0001 to 10 wt%.
- 25 30. An insecticidal composition according to claim 28 or claim 29, in which the preservative is selected from sorbic acid and salts thereof, methylparaben, propylparaben and mixtures thereof.
- 30 31. An insecticidal composition according to any preceding claim further comprising an anti-oxidant.
32. An insecticidal composition according to claim 31, in which the anti-oxidant is present in an amount of from 0.0001 to 10 wt%.
- 35 33. An insecticidal composition according to claim 31 or claim 32, in which the anti-oxidant is selected from TBHQ, butylated hydroxytoluene, butylated hydroxyanisole and mixtures thereof.

34. An insecticidal composition according to any preceding claim further comprising a human taste deterrent
- 5 35. An insecticidal composition according to claim 34, in which the human taste deterrent is present in an amount of from 1 to 200 ppm.
- 10 36. An insecticidal composition according to claim 34 or claim 35, in which the human taste deterrent is denatonium benzoate.
- 15 37. An insecticide composition according to any preceding claim further comprising a colorant.
- 20 38. An insecticidal composition according to any preceding claim further comprising one or more other components selected from feeding stimulants, and fillers and animal taste deterrents.
- 25 39. An insecticidal composition according to any preceding claim having a penetration hardness of from 0.1 to 10 mm.
40. An insecticidal composition according to any preceding claim having a penetration hardness of from 0.5 to 5 mm.
- 30 41. An insecticidal composition according to any preceding claim having a melting point of at least 20 °C, preferably at least 35 °C, more preferably at least 40 °C and most preferably at least 45 °C.
- 35 42. An insecticidal composition according to any preceding claim having a setting point in the range

from 20 to 200°C, preferably from 30 to 70°C and more preferably from 45-55 °C.

43. A method for controlling insects in which a solid
5 stick insecticidal composition according to any one
of claims 1 to 42 is applied and deposited on to a
hard surface to be treated.
44. A method for controlling insects according to claim
10 43, in which the insecticidal composition is applied
as a line of material by drawing the stick along the
surface.
45. A method for controlling insects according to claim
15 43, in which the insecticidal composition is applied
as a series of spots on the surface.
46. A method for controlling insects according to claim
20 43, in which the insecticidal composition is applied
as a smear on the surface.
47. A method for controlling insects according to any one
of claims 43 to 46, in which the insects which crawl.
- 25 48. A method for controlling insects according to claim
47, in which the insects are cockroaches.
49. An insect control product comprising a solid stick
insecticidal composition according to any one of
30 claims 1 to 42 and a package.
50. An insect control product according to claim 49 in
which the package comprises a protective wrapping.

51. An insect control product according to claim 50 in which the protective wrapping comprises paper, plastic, fabric, metal foil or laminates.
- 5 52. An insect control product according to claim 50 or claim 51, in which the wrapping includes tear lines or points of weakness at intervals along the length of the wrapping thus allowing for progressive removal of circumferential strips of wrapping from the stick.
- 10 53. An insect control product according to claim 49 in which the package comprises an applicator including a bottom, in which is held one end of the stick, and a tubular lid, which partially or fully encloses the
- 15 insecticidal stick.
54. An insect control product according to claim 49 in which the package comprises an applicator in which the insecticidal stick is extendable from a casing.
- 20 55. An insect control product according to any one of claims 49 to 54 in which the package comprises a label or a label is applied to the package.
- 25 56. Use of an insecticide and a food material in the manufacture a solid stick insecticidal composition according to any one of claims 1 to 42 for application to a hard surface to be treated.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2004/003658

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A01N25/00 A01N25/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 162 575 A (LANG DICKSON S) 22 December 1964 (1964-12-22) cited in the application the whole document	1-56
X	DATABASE WPI Section Ch, Week 197947 Derwent Publications Ltd., London, GB; Class C03, AN 1979-84995B XP002309864 & JP 54 132237 A (TAISHO PHARM CO LTD) 15 October 1979 (1979-10-15) abstract	1-56
A	GB 2 221 392 A (SHAO CHIA MU) 7 February 1990 (1990-02-07)	

-/-

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

9 December 2004

Date of mailing of the international search report

30/12/2004

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Fort, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2004/003658

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 473 582 A (GREENE C LAWRENCE) 25 September 1984 (1984-09-25) cited in the application -----	
A	US 3 826 232 A (DUFFEY T ET AL) 30 July 1974 (1974-07-30) cited in the application -----	
A	WO 95/17814 A (MINNESOTA MINING & MFG) 6 July 1995 (1995-07-06) -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB2004/003658

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3162575	A	22-12-1964	NONE	
JP 54132237	A	15-10-1979	NONE	
GB 2221392	A	07-02-1990	NONE	
US 4473582	A	25-09-1984	EP 0046336 A2	24-02-1982
US 3826232	A	30-07-1974	NONE	
WO 9517814	A	06-07-1995	AU 1215395 A	17-07-1995
			CA 2178078 A1	06-07-1995
			EP 0737034 A1	16-10-1996
			JP 9507227 T	22-07-1997
			NZ 277205 A	26-02-1998
			WO 9517814 A1	06-07-1995
			US 5626859 A	06-05-1997
			ZA 9410286 A	24-06-1996